

On analytic descriptions of two-dimensional surfaces associated with the $\mathbb{C}P^{N-1}$ sigma model

A. M. Grundland^{1,2 *} and İ. Yurduşen^{1 †}

¹Centre de Recherches Mathématiques, Université de Montréal,
CP 6128, Succ. Centre-Ville, Montréal, Québec H3C 3J7, Canada

²Université du Québec, Trois-Rivières, CP500, QC, G9A 5H7, Canada

May 15, 2009

Abstract

We study analytic descriptions of conformal immersions of the Riemann sphere S^2 into the $\mathbb{C}P^{N-1}$ sigma model. In particular, an explicit expression for two-dimensional (2-D) surfaces, obtained from the generalized Weierstrass formula, is given. It is also demonstrated that these surfaces coincide with the ones obtained from the Sym-Tafel formula. These two approaches correspond to parametrizations of one and the same surface in $\mathbb{R}^{N^2-1}$.

Key words: Sigma models, Weierstrass formula for immersion, Sym-Tafel formula

PACS numbers: 02.40.Hw, 02.20.Sv, 02.30.Ik

In this note we investigate the relations between the $\mathbb{C}P^{N-1}$ sigma model and the generalized Weierstrass formula for immersion of 2-D surfaces in multi-dimensional Euclidean spaces. These links have been discussed in [1, 2, 3] and are governed by the formula

$$X_k(\xi, \bar{\xi}) = i \int_{\gamma} (-[\partial P_k, P_k] d\xi + [\bar{\partial} P_k, P_k] d\bar{\xi}), \quad k = 0, 1, \dots, N-2, \quad (1)$$

where P_k are rank-1 orthogonal projectors which satisfy the completely integrable 2-D $\mathbb{C}P^{N-1}$ sigma model

$$\partial[\bar{\partial} P_k, P_k] + \bar{\partial}[\partial P_k, P_k] = 0, \quad P_k^2 = P_k, \quad P_k^\dagger = P_k. \quad (2)$$

*E-mail address: grundlan@crm.umontreal.ca

†E-mail address: yurdusen@crm.umontreal.ca

We first demonstrate that for any solution of the $\mathbb{C}P^{N-1}$ sigma model (2) defined on the Riemann sphere S^2 with a finite action functional, the generalized Weierstrass formula for immersion of 2-D surfaces (1) can be integrated explicitly up to a constant of integration and expressed in terms of the projectors P_k

$$X_k(\xi, \bar{\xi}) = -i(P_k + 2 \sum_{j=0}^{k-1} P_j), \quad k = 0, 1, \dots, N-2. \quad (3)$$

Indeed if we assume that the $\mathbb{C}P^{N-1}$ sigma model is defined on the sphere S^2 with a finite action functional, then the complete set of regular solutions are known [4, 5]. As a result, one gets three classes of solutions, namely (i) holomorphic (i.e. $\bar{\partial}f = 0$), (ii) antiholomorphic (i.e. $\partial f = 0$) and (iii) mixed. The mixed solutions can be determined from either the holomorphic or the antiholomorphic nonconstant functions by the successive application of the operator P_+ [6]

$$P_+ : f \in \mathbb{C}^N \rightarrow P_+ f = \partial f - f \frac{f^\dagger \partial f}{f^\dagger f}, \quad (4)$$

where f is any nonconstant holomorphic function. This allows one to construct mixed solutions $f_k = P_+^k f$ which represent harmonic maps from S^2 to the $\mathbb{C}P^{N-1}$ model. Here, the operator P_+^k is obtained by applying successively k times the operator P_+ and $P_+^0 = \text{id}$. Hence, using (4) for every $k \leq N-1$ we can define a set of rank-1 projectors $\{P_0, P_1, \dots, P_k\}$

$$P_k := \frac{f_k \otimes f_k^\dagger}{f_k^\dagger \cdot f_k}, \quad k = 0, 1, \dots, N-1, \quad (5)$$

which determine conservation laws of the form (2). The first ($k = 0$) and the last ($k = N-1$) conservation laws are related to the holomorphic and antiholomorphic solutions, respectively, while the intermediate ones are related to the mixed solutions. Consequently, according to the Weierstrass procedure we can obtain 2-D surfaces for each projector P_k . By straightforward calculation one gets [2]

$$\begin{aligned} [\partial P_k, P_k] &= \partial P_k + 2 \frac{(P_+^k f) \otimes (P_+^{k-1} f)^\dagger}{|P_+^{k-1} f|^2}, \\ [\bar{\partial} P_k, P_k] &= -\bar{\partial} P_k - 2 \frac{(P_+^{k-1} f) \otimes (P_+^k f)^\dagger}{|P_+^{k-1} f|^2}. \end{aligned} \quad (6)$$

Hence, for every $k \leq N-1$, the Weierstrass formula for immersion (1) takes the form

$$dX_k = -i \left[(\partial P_k + 2 \frac{(P_+^k f) \otimes (P_+^{k-1} f)^\dagger}{|P_+^{k-1} f|^2}) d\xi + (\bar{\partial} P_k + 2 \frac{(P_+^{k-1} f) \otimes (P_+^k f)^\dagger}{|P_+^{k-1} f|^2}) d\bar{\xi} \right]. \quad (7)$$

Note that the two surfaces corresponding to $k = 1$ and $k = N-1$ are precisely the same objects, since one gets antiholomorphic solutions of the model after applying $N-1$ times the operator P_+ to the nonconstant holomorphic function

f . Hence, there appear at most $N - 2$ different surfaces as a result of the generalized Weierstrass formula.

The integration of (7) gives us (3) which can be shown as follows: It is immediately seen that for $k = 0$ we have $X_0 = -iP_0$ and upon differentiation we obtain

$$dX_0 = -i[\partial P_0 d\xi + \bar{\partial} P_0 d\bar{\xi}], \quad (8)$$

which coincides with (7) for $k = 0$. For $k = 1$, we need to show that

$$\partial P_0 = \frac{(P_+ f) \otimes f^\dagger}{|f|^2}, \quad (9)$$

which could easily be computed by differentiating P_0 and bearing in mind that f is holomorphic. In order to show that (3) holds for any k we assume that

$$\partial(P_0 + P_1 + \cdots, P_{k-2}) = \frac{(P_+^{k-1} f) \otimes (P_+^{k-2} f)^\dagger}{|P_+^{k-2} f|^2}, \quad (10)$$

and then compute ∂P_{k-1}

$$\begin{aligned} \partial P_{k-1} &= \partial \left[\frac{(P_+^{k-1} f) \otimes (P_+^{k-1} f)^\dagger}{|P_+^{k-1} f|^2} \right] \\ &= \frac{(P_+^k f) \otimes (P_+^{k-1} f)^\dagger}{|P_+^{k-1} f|^2} - \frac{(P_+^{k-1} f) \otimes (P_+^{k-2} f)^\dagger}{|P_+^{k-2} f|^2}, \end{aligned} \quad (11)$$

where we have used the fact that

$$\partial(P_+^{k-1} f)^\dagger = -\frac{|P_+^{k-1} f|^2}{|P_+^{k-2} f|^2} (P_+^{k-2} f)^\dagger, \quad (12)$$

together with the orthogonality relation

$$(P_+^k f)^\dagger \cdot (P_+^l f) = 0, \quad \text{for } k \neq l. \quad (13)$$

Thus, we have shown that

$$\partial \left(\sum_{j=0}^{k-1} P_j \right) = \frac{(P_+^k f) \otimes (P_+^{k-1} f)^\dagger}{|P_+^{k-1} f|^2}, \quad (14)$$

which indeed justifies (3).

It is interesting to note that this immersion function X_k coincides with the results obtained in [2], namely the surface for nonholomorphic Veronese type solutions lives in $\mathbb{R}^3$.

For the $\mathbb{C}P^{N-1}$ sigma model, it may also be of interest to investigate the links between the generalized Weierstrass formula for the immersion of 2-D surfaces in the $su(N)$ algebra and the approach based on the linear spectral problem for constructing infinitely many surfaces in multi-dimensional Euclidean spaces $\mathbb{R}^{N^2-1} \simeq su(N)$.

Following the approach proposed by A. Sym and J. Tafel in [7, 8, 9, 10], in particular using their formula

$$X_k(\xi, \bar{\xi}) = \alpha(\lambda) \phi_k^{-1} \frac{\partial \phi_k}{\partial \lambda}, \quad \phi_k \in SU(N), \quad k = 0, 1, \dots, N-2, \quad (15)$$

for integrable surfaces derived via the Lax pair [11, 12]

$$\partial \phi_k = \frac{2}{1+\lambda} [\partial P_k, P_k] \phi_k, \quad \bar{\partial} \phi_k = \frac{2}{1-\lambda} [\bar{\partial} P_k, P_k] \phi_k, \quad k = 0, 1, \dots, N-1, \quad (16)$$

we demonstrate that there exist 2-D surfaces with $su(N)$ -valued immersion functions $X_k(\xi, \bar{\xi})$ which are precisely of the form (3). Here, λ is a spectral parameter and the compatibility conditions for the system (16) coincide with the $\mathbb{C}P^{N-1}$ model equations (2). Furthermore, $\alpha(\lambda)$ is some scalar function of λ .

In the purely instantonic case (i.e. holomorphic and antiholomorphic solutions), the orthogonal projector P has the form

$$P_0 = \frac{f \otimes f^\dagger}{f^\dagger \cdot f}, \quad (17)$$

which satisfies

$$[\partial P_0, P_0] = \frac{P_+ f \otimes f^\dagger}{f^\dagger \cdot f}, \quad [\bar{\partial} P_0, P_0] = -\frac{f \otimes (P_+ f)^\dagger}{f^\dagger \cdot f}. \quad (18)$$

Looking for a solution $\phi_0 = \phi_0(\lambda)$ of the linear spectral problem (16) when ϕ_0 tends to 1 as $\lambda \rightarrow \infty$, we make the Ansatz [6]

$$\phi_0 = I_N - \frac{2}{1-\lambda} P_0, \quad (19)$$

where I_N is the $N \times N$ identity matrix. The inverse matrix of ϕ_0 is given by

$$\phi_0^{-1} = I_N - \frac{2}{1+\lambda} P_0. \quad (20)$$

Hence, according to the Sym-Tafel formula (15), the surface associated with the $\mathbb{C}P^{N-1}$ model is given up to an additive $su(N)$ -valued constant by

$$X_0 = \frac{2}{1-\lambda^2} P_0. \quad (21)$$

For the nonholomorphic solutions (i.e. the mixed solutions), we proceed in an analogous way and find that [6],

$$\phi_k = I_N + \frac{4\lambda}{(1-\lambda)^2} \sum_{j=0}^{k-1} P_j - \frac{2}{1-\lambda} P_k, \quad k = 1, \dots, N-2, \quad (22)$$

and the inverse of ϕ_k has the form

$$\phi_k^{-1} = I_N - \frac{4\lambda}{(1+\lambda)^2} \sum_{j=0}^{k-1} P_j - \frac{2}{1+\lambda} P_k. \quad (23)$$

The Sym-Tafel formula (15) for the immersion function $X_k(\xi, \bar{\xi})$ of 2-D surfaces associated with the $\mathbb{C}P^{N-1}$ model is given up to an additive $su(N)$ -valued constant by the formula

$$X_k(\xi, \bar{\xi}) = \frac{2}{1 - \lambda^2} \left(P_k + 2 \sum_{j=0}^{k-1} P_j \right), \quad k = 0, 1, \dots, N-2. \quad (24)$$

This result for immersion of 2-D surfaces in the $su(N)$ Lie algebras coincides with the one obtained from the Weierstrass representation (3) and shows the equivalence between these two approaches.

To conclude, we give an explicit expression for 2-D surfaces, obtained from the generalized Weierstrass formula and demonstrate that these surfaces coincide with the ones obtained from the Sym-Tafel formula.

Acknowledgments

This work is supported in part by research grants from NSERC of Canada. İ.Y. acknowledges a postdoctoral fellowship awarded by the Laboratory of Mathematical Physics of the CRM, Université de Montréal.

References

- [1] A. M Grundland, W. A. Hereman, and İ. Yurduşen 2008 Conformally parametrized surfaces associated with $\mathbb{C}P^{N-1}$ sigma models J. Phys. A: Math. Theor. **41** 065204
- [2] A. M Grundland and İ. Yurduşen 2008 Surfaces obtained from $\mathbb{C}P^{N-1}$ sigma models Int. J. Mod. Phys. A **23** 5137-5157
- [3] A. M Grundland, A. Strasburger, and W. J. Zakrzewski 2006 Surfaces immersed in $su(N+1)$ Lie algebras obtained from the $\mathbb{C}P^N$ sigma models J. Phys. A: Math. Gen. **39** 9187-9113
- [4] A. M. Din and W. J. Zakrzewski 1980 General classical solutions of the $\mathbb{C}P^{N-1}$ model Nucl. Phys. **B174** 397-403
- [5] R. Sasaki 1983 General classical-solutions of the complex Grassmannian and $\mathbb{C}P^{N-1}$ sigma models Phys. Lett. **B130** 69-72
- [6] W. J. Zakrzewski 1989 *Low Dimensional Sigma Models* (Bristol: Adam Hilger)
- [7] A. Sym 1982 Soliton surfaces Lett. Nuovo Cimen. **33** 394-400
- [8] A. Sym 1983 Soliton surfaces II. Geometric unification of solvable nonlinearities Lett. Nuovo Cimen. **36** 307-312
- [9] J. Tafel 1995 Surfaces in $\mathbb{R}^3$ with prescribed curvature J. Geom. Phys. **17** 381-390

- [10] A. Sym 1985 Soliton surfaces and their applications (Soliton geometry from spectral problems), in: *Geometric Aspects of the Einstein Equations and Integrable Systems*, Lecture Notes in Physics No: **239** Ed.: R. Martini (Berlin: Springer-Verlag)
- [11] A. V. Mikhailov 1986 Integrable magnetic models, in: *Solitons*, Eds.: S. E. Trullinger, V. E. Zakharov, and V. L. Pokrovsky (Modern Problems in Condensed Matter, Vol. 17) (Amsterdam: North-Holland) pp. 623–690.
- [12] V. E. Zakharov and A. V. Mikhailov 1979 Relativistically invariant two-dimensional models of field theory which are integrable by means of the inverse scattering problem method Sov. Phys. **40** 1017-1049